



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2007

Measuring dark energy with the shear triplet statistics

Sereno, M

Abstract: The shear triplet statistics is a geometric method to measure cosmological parameters with observations in the weak gravitational lensing regime towards massive haloes. Here, this proposal is considered to probe the dark energy equation of state and its time derivative in view of future wide-field galaxy surveys. A survey with a median redshift of 0.7 and a total area of 10 000 deg² would be pretty effective in determining the dark matter cosmological density and in putting useful constraints on the dark energy properties

DOI: <https://doi.org/10.1111/j.1365-2966.2007.11600.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-155284>

Journal Article

Published Version

Originally published at:

Sereno, M (2007). Measuring dark energy with the shear triplet statistics. *Monthly Notices of the Royal Astronomical Society*, 377(1):229-233.

DOI: <https://doi.org/10.1111/j.1365-2966.2007.11600.x>

Measuring dark energy with the shear triplet statistics

M. Sereno[★]

Institut für Theoretische Physik, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

Accepted 2007 February 2. Received 2007 February 1; in original form 2006 December 15

ABSTRACT

The shear triplet statistics is a geometric method to measure cosmological parameters with observations in the weak gravitational lensing regime towards massive haloes. Here, this proposal is considered to probe the dark energy equation of state and its time derivative in view of future wide-field galaxy surveys. A survey with a median redshift of ~ 0.7 and a total area of $\sim 10\,000$ deg² would be pretty effective in determining the dark matter cosmological density and in putting useful constraints on the dark energy properties.

Key words: gravitational lensing – cosmology: observations – large-scale structure of Universe.

1 INTRODUCTION

The deep theoretical understanding of gravitational lensing makes it an attractive probe of dark energy, one of the main puzzles of modern cosmology. Dark energy can show up either in a pure geometrical way by affecting the distance redshift relation or via its effect on the growth of structure. Lensing is sensitive to both signatures since a ratio of distances appears in the scaling of lensing parameters with redshifts and because the effective mass of lensing structures reflects the power spectrum and growth rate of large-scale density perturbations (Bartelmann & Schneider 2001; Munshi et al. 2006, and references therein). Probing dark energy through both geometry and growth by taking weak gravitational lensing two-point functions (such as the shear power spectrum) of distant galaxy images as a function of redshift can be appealing (Huterer 2002; Song & Knox 2004; Takada & Jain 2004; Heavens, Kitching & Taylor 2006, and references therein), but this method relies on the interpretation of the distortion signal on scales where non-linear evolution necessitates accurate modelling from numerical simulations. In the strong lensing regime, the distance ratio and the cosmological parameters can be derived using observations of giant luminous arcs in rich clusters of galaxies but the mass profile and the mass normalization of the lenses must be assumed to be known, for example, from X-ray measurements (Sereno 2002; Sereno & Longo 2004, and references therein). Strong gravitational lensing observations of image separations and time delays could provide useful constraints as well (Linder 2004).

The basic idea behind geometric methods is to separate out the information purely from the distance ratios, irrespective of the lensing mass distribution. In the strong lensing regime, this can be achieved by considering multiple image systems and the ratios of arc angular positions, but this still requires the assumption that the lensing mass, even if unknown, is pretty regular (Golse, Kneib & Soucail

2002). The weak lensing regime seems more promising. Jain & Taylor (2003) proposed to take the ratio of the galaxy–shear correlation function at different redshifts behind galaxy groups and galaxy clusters. Dark energy parameter constraints based on this geometric method, which extends into the non-linear matter power spectrum but still drops out the mass of the lens, have been then discussed by several authors in view of future galaxy surveys (Bernstein & Jain 2004; Hu & Jain 2004; Song & Knox 2004). The method was then further developed by directly considering ratios of the shears behind a cluster without first generating the cross-correlation functions (Taylor et al. 2007). Alternatively, Zhang, Hui & Stebbins (2005) pointed out how the cross-correlation of a foreground galaxy-density field or shear field with the shear field from a background source population scales with the source redshift in a way that can be used to constrain cosmology without making assumptions about the mass/galaxy power spectrum.

In this paper, we want to reconsider the so-called triplet statistics, an original idea to constrain cosmological parameters from weak lensing in galaxy clusters proposed in Gautret, Fort & Mellier (2000). This method is able to disentangle the effect of the lensing mass, described by local convergence and shear terms, from the cosmological parameters by considering the ellipticities in triplets of galaxies located at about the same angular position but having different redshifts. Differently from similar proposals (Taylor et al. 2007), the triplet statistics are not limited to the outskirts of massive haloes and can be used even in the inner regions. Gautret et al. (2000) originally considered the method for determining the cosmological constant with observations towards very massive galaxy clusters. Here, we review the triplet statistics and discuss the measurement of the dark energy properties with future weak lensing survey.

2 BASICS

The distortion of images of background galaxies is determined by the convergence k , i.e. the lensing strength, and the complex shear $\gamma = \gamma_1 + i\gamma_2$. The lensing parameters can be related to the value they

[★]E-mail: sereno@physik.unizh.ch

would have for a source at a reference redshift (Seitz & Schneider 1997; Gautret et al. 2000)

$$k = \omega(z_s)k_{\text{ref}}, \quad (1)$$

$$\gamma = \omega(z_s)\gamma_{\text{ref}}. \quad (2)$$

The lensing factor $\omega(z_s)$ is defined as a ratio of angular diameter distances and contains the cosmological dependency,

$$\omega(z_s) \equiv \frac{D_{\text{ds}}}{D_{\text{os}}} \left(\frac{D_{\text{ds}}}{D_{\text{os}}} \Big|_{z_s=z_{\text{ref}}} \right)^{-1}, \quad (3)$$

where D_{ij} is the angular diameter distance between the redshifts z_i and z_j with the redshifts of interest being those of the observer o , the lens d and the source s . We consider a standard Friedmann–Lemaître–Robertson–Walker model of universe, filled with non-interacting pressureless matter and dark energy, parametrized through its equation of state $w \simeq w_o + w_a(1-a)$, with $a \equiv 1/(1+z)$. Since the contribution from relativistic particles is negligible in the redshift range investigated in our analysis, we will neglect it in what follows. In such a model of universe, the angular diameter distance between an observer at z_i and a source at z_j is

$$D_{ij} = \frac{c}{H_0} \frac{1}{1+z_j} \frac{1}{|\Omega_{\text{K}0}|^{1/2}} \text{Sinn} \left[\int_{z_i}^{z_j} \frac{|\Omega_{\text{K}0}|^{1/2}}{H(z)/H_0} dz \right], \quad (4)$$

where

$$\frac{H(z)}{H_0} = \sqrt{\Omega_{\text{M}}a^{-3} + \Omega_{\text{v}}a^{-3(1+w_0+w_a)}e^{-3w_a(1-a)} + \Omega_{\text{K}}a^{-2}}, \quad (5)$$

and H_0 is the present value of the Hubble parameter; Ω_{M} and Ω_{v} are the today normalized densities of dust and dark energy, respectively; $\Omega_{\text{K}} \equiv 1 - \Omega_{\text{M}} - \Omega_{\text{v}}$; Sinn is defined as being \sinh when $\Omega_{\text{K}} > 0$, \sin when $\Omega_{\text{K}} < 0$ and as the identity when $\Omega_{\text{K}} = 0$. For the expression of the distance in an inhomogeneous universe, we refer to Sereno et al. (2001) and Sereno, Piedipalumbo & Sazhin (2002).

The transformation from intrinsic to observed ellipticity in the weak lensing regime ($\gamma, k \ll 1$) takes a simple form (Seitz & Schneider 1997; Gautret et al. 2000). Due to a lensing halo, a galaxy with intrinsic ellipticity ϵ_s is imaged with ellipticity (Gautret et al. 2000)

$$\epsilon \simeq (1 - g^2)\epsilon_s + \mathbf{g} \quad (6)$$

with $\mathbf{g} \equiv \gamma/(1-k)$ being the complex reduced shear. Background galaxies at (nearly) the same angular position probe the same local cluster mass distribution and gravitational potential. Then, comparing the shear amplitude for three galaxies having different redshift allows to separate the effect of the mass distribution from cosmology. A geometrical operator can be built from the measured ellipticities ϵ_i and redshifts z_i of the three galaxies $i = \{a, b, c\}$ in such a way that it depends only on cosmology (Gautret et al. 2000)

$$T_{abc} = \begin{pmatrix} 1 & \omega_a & \omega_a \epsilon_b \epsilon_c^* \\ 1 & \omega_b & \omega_b \epsilon_c \epsilon_a^* \\ 1 & \omega_c & \omega_c \epsilon_a \epsilon_b^* \end{pmatrix}. \quad (7)$$

Redshifts of the galaxies inside triplets are in ascending order. For three intrinsically spherical galaxies, $T_{abc} = 0$ when the lensing factors ω_i are computed for the actual values of the cosmological parameters, apart from noise. T_{abc} is linear regarding the ellipticities, which makes the principal source of noise randomly distributed around zero. Apart from noise, the main part of the triplet operator

contains the cosmological dependence and can be approximated as

$$T_{abc}^m \simeq \begin{vmatrix} 1 & \omega_a & \omega_a/\omega_a^0 \\ 1 & \omega_b & \omega_b/\omega_b^0 \\ 1 & \omega_c & \omega_c/\omega_c^0 \end{vmatrix} \omega_a^0 \omega_b^0 \omega_c^0 \gamma_{\text{ref}}^2, \quad (8)$$

where the apex 0 denotes that the angular diameter distances have been calculated for the actual values of the cosmological parameters and where we have considered only one component of the shear. If we neglect the contribution from the local convergence, the reduced shear \mathbf{g} can be identified with the local shear and then it is enough to take the ratio of the observed ellipticities of a pair of near galaxies in order to separate the effect of cosmology (Taylor et al. 2007). Neglecting k introduces a systematic error that can be significant for the largest clusters. A 10 per cent variation of the cosmological parameters changes the reduced shear by about 1 per cent which is 10 times smaller than the relative variation due to the $1 - \omega k_{\text{ref}}$ for $k_{\text{ref}} \lesssim 0.1$ (Gautret et al. 2000). Even if the majority of the signal comes from intermediate-mass clusters, the highest return in accuracy comes from the largest haloes (Taylor et al. 2007) so that this effect must be properly accounted for.

Let us sort the redshift in a triplet such that $z_a < z_b < z_c$. From the matrix form of the operator T_{abc} , it is clear that if two galaxies in a triplet are very close, then $T_{abc} \simeq 0$ with no regard to the cosmological parameters. $T_{abc} \simeq 0$ also if the minimum redshift in the triplet is very close to the lens redshift. Once fixed the minimum and the maximum redshift in a triplet, the sensitivity of the operator T_{abc} is maximized for an intermediate redshift z_b nearly in the middle of the redshift range. Once fixed z_a and z_b , the sensitivity increases with the maximum redshift z_c . Hence, the main information from the triplet method comes from the high-redshift tail of the background source distribution.

3 FORECAST FOR LENSING SURVEYS

There are a number of current and planned imaging surveys for weak lensing analyses (Peacock et al. 2006). Beyond 2007, the funded VST (VLT Survey telescope) public survey The Kilo-Degree Survey (KIDS) will cover at least 1500 deg² in four broad-band optical filters. Combined with a following near-infrared coverage by the Visible and Infrared Survey Telescope for Astronomy (VISTA), this will yield a nine-band optical-IR survey with depth approximately 2 mag deeper than the Sloan and with very accurate photometric redshift estimate. The typical KIDS lenses will be at $z \sim 0.2-0.4$. Beyond KIDS, the planned Dark Energy Survey on the CTIO Blanco telescope or the dark CAM survey at VISTA is expected to take another steps forward in terms of sky coverage, imaging $\sim 10\,000$ deg². In what follows, we give a cosmological parameter estimation forecast for such surveys.

The background redshift distribution for a typical magnitude-limited survey can be taken to be

$$\frac{dn_{\text{gal}}}{dz} = n_0 \frac{3}{2} \frac{z^2}{z_0^3} \exp \left[- \left(\frac{z}{z_0} \right)^{3/2} \right]. \quad (9)$$

The redshift scale z_0 is related to the median redshift of the survey, z_m , by $z_0 = z_m/1.412$; n_0 gives the total number density of sources with usable photometric redshift and shape estimate. For a KIDS-like survey, we can take $z_m \sim 0.7$.

The main part of the lensing signal comes from halo lensing masses in the range $5 \times 10^{13} \lesssim M/M_{\odot} \lesssim 10^{15}$. The number density of haloes can be accurately calculated and varies from $n_{\text{hal}} \gtrsim 10^2$ per square degree for $M \sim 10^{13}$ to $n_{\text{hal}} \gtrsim 10^{-2}$ per square degree for

$M \sim 10^{15}$ (Taylor et al. 2007). An aperture size $\theta_{\max} \simeq 2\text{--}5$ arcmin corresponds roughly to the virial radii of such massive haloes over the redshift range considered. An inner circle with aperture $\theta_{\text{in}} \simeq 0.1\text{--}0.8$ arcmin has to be excised from the data to exclude the arclets and the strong lensing regime. Considering a simple isothermal sphere model for the lenses, we get that $\langle \gamma_{\text{ref}} \rangle \sim 0.04\text{--}0.15$ in the mass range of interest. We remark as the triplet operator is proportional to $\langle \gamma_{\text{ref}}^2 \rangle$ rather than $\langle \gamma_{\text{ref}} \rangle^2$. These estimates for the shear are lower limits since considering a Navarro–Frank–White profile as a deflector, the shear signal would increase for the same halo mass.

In order to study the sensitivity of the triplet method on the cosmological parameters, we perform a χ^2 analysis. The main difference with the statistical analysis in Gautret et al. (2000) is that they considered a mean triplet operator as the average over all triplets whereas here we consider a χ^2 built on the linearly independent triplets. Not all of the triplets we can put together behind a lens contain independent information. It can be easily shown that from N_g near galaxies only $N_g - 2$ triplets out of the binomial factor $(N_g, 3)$ are independent. If a triplet contains at least one galaxy not already included in the sample of other triplets, then it is linearly independent. Triplet selection can be properly optimised. As maximum distance between the triplet components, we take a separation of $\Delta r \sim 20$ arcsec (Gautret et al. 2000). This is a good balance between having a sufficient number of triplets and not smearing the signal. As χ^2 , we consider

$$\chi^2 = \sum_{l,(a,b,c)} \left(\frac{T_{abc}^m}{\delta T_{abc}} \right)^2 \quad (10)$$

with the sum running over the lensing haloes l optically selected in the survey and the independent triplets for each halo. Foreground lenses and background sources are modelled according to the previous discussion. For each background galaxy, we completed the triplet by selecting the two neighbouring galaxies ($\Delta r \lesssim 20$ arcsec) which maximize the signal.

Statistical and systematic errors affecting the method have been deeply discussed in Gautret et al. (2000). The main sources of noise are (i) the intrinsic source ellipticities; (ii) the errors on measured ellipticities; (iii) the fact that sources do not experience exactly the same potential and finally (iv) the errors on measured (photometric) redshifts. It can be shown that due to the linearity of the operator, the noise is linear with respect to each individual term and then is proportional to $1/\sqrt{N}$ with N the total number of triplets. The dominant contribution to the error budget comes from the intrinsic ellipticity.

Together with statistical noise, several systematics affect the method. The main ones are well understood and have been identified as (i) a bias due to an asymmetry in the probability distribution of the terms $\Delta\omega$ due to photometric redshift errors and, mainly, (ii) contamination by background structure, either galaxy–galaxy lensing or large-scale structure. Sources in very close angular pairs, for which another galaxy could play the role of lens, could be rejected from the sample. It can be shown that for surveys large enough, the effect of large-scale structure can be neglected at first order with respect to the statistical noise (Gautret et al. 2000; Taylor et al. 2007). The dominant noise source is due to intrinsic ellipticity dispersion (Gautret et al. 2000; Taylor et al. 2007). Then, as an estimate of δT_{abc} in equation (10) we will take

$$\delta T_{abc}^2 = \frac{\sigma_\epsilon^2}{2} \sum_{i=a,b,c} \left(\frac{\partial T_{abc}}{\partial \epsilon_i} \right)^2 \Big|_{\epsilon_i = \omega_i^0 \gamma_{\text{ref}}} \quad (11)$$

where the factor of 2 in the denominator arises because we are using




Figure 1. Confidence regions in the $\Omega_M - \Omega_\Lambda$ plane as expected for a lensing survey with $z_m \sim 0.7$. Contours show the 1, 2, 3 σ limits around the fiducial model $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$. The inner thin and the outer thick contours refer to a survey with total area $A = 10\,000$ deg² and 1500 deg², respectively.

only one component of the measured ellipticity. As intrinsic ellipticity dispersion, we take $\sigma_\epsilon = 0.3$. The effect of lensing by large-scale structure will be discussed later. However, even if we are underestimating the statistical noise by considering only the main contributor, the choice of parameters has been generally conservative. Results for a survey with median redshift ~ 0.7 and a typical density of $n_0 \sim 30$ galaxies per square arcmin are shown in Figs 1–3. As a first




Figure 2. Confidence regions in the $\Omega_M - w_0$ plane for a lensing survey covering 10 000 deg². Contours show the 1, 2, 3 σ limits around the fiducial flat model $\Omega_M = 0$, $w_0 = -1$ and assuming no evolution for the dark energy, $w_a = 0$. The dashed line separates models with either accelerated or decelerated expansion.




Figure 3. Confidence regions in the $w_0 - w_a$ plane for a lensing survey covering $10\,000 \text{ deg}^2$. Contours show the 1, 2, 3σ limits around a reference flat model $w_0 = -1$, $w_a = 0$ assuming flat geometry and a sharp prior $\Omega_M = 0.3$.

step, we have considered dark energy in the form of a cosmological constant ($w_0 = -1$ and $w_a = 0$; see Fig. 1). The contours show the 1, 2 and 3σ confidence limits for two parameters ($\Delta\chi^2 = 2.30$, 6.16 and 11.8, respectively). With a survey area of $10\,000 \text{ deg}^2$, the triplet statistics can constrain the matter density parameter but is pretty insensitive to the total amount of vacuum energy. This degeneracy changes with redshift. For tests in the strong lensing regime based on luminous giant arcs, the contours are nearly orthogonal to the triplet estimator since the distance ratio is evaluated for very high redshift sources (Sereno & Longo 2004).

Assuming a flat model and a constant equation of state, the triplet statistics can put a quite firm upper limit on w_0 (see Fig. 2). On the other hand, the confidence regions spread out well within the phantom regime ($w_0 < -1$), as usual for methods based on the distance–redshift relation. The phantom regime would be also consistent with a higher value of the matter density parameter. Contours in the $w_0 - w_a$ plane are still pretty elongated (see Fig. 3). On its own, the triplet statistics cannot say much about the evolution with time of the dark energy, but if combined with orthogonal methods such as the cosmic microwave background radiation or the baryonic acoustic oscillation in the matter power spectrum, constraints could be significant.

The linear size of the confidence regions shrinks approximately as $1/\sqrt{N_g}$, with N_g the total number of galaxies in the survey. Cosmological constraints would then strongly benefit by a very large survey area. Furthermore, increasing the median redshift of the survey would both increase the local galaxy density and probe the distance ratio in a redshift range more sensitive to cosmology.

Together with the variance term proportional to the intrinsic uncertainty per shear mode due to the galaxy intrinsic ellipticities (σ_γ), which is related to the shape noise and to the shot noise, the other main source of error is due to lensing by large-scale structure in between the lens and the sources (σ_{LLS}), related to the sampling variance term (Hu & Jain 2004; Zhang et al. 2005). Depending on the survey strategy, these two terms can be comparable. Taylor et al.

(2007) gave a simple approximate scaling relation between the two terms for a survey of median redshift z_m , $\sigma_{\text{LLS}}^2 \simeq (24.1 z_m^4 \Delta z) \sigma_\gamma^2$, where Δz is the typical photometric redshift error for the survey. Then, for $z_m \sim 0.7$ and $\Delta z \sim 0.05$, error estimates on cosmological parameters obtained considering just the shot noise should be increased by ~ 15 per cent.

4 FINAL REMARKS

The aim of this paper has been the evaluation of the triplet statistics as a dark energy probe in view of future galaxy survey. The main source of statistical uncertainty in our statistical approach was the intrinsic source ellipticity. As regards the main systematics, the contribution of large-scale structure to the observed shear, being uncorrelated with the cluster effect, should average out over independent clusters along different lines of sight (Taylor et al. 2007). In a survey measuring photometric redshifts, data should be collected in redshift bins with width equal to the typical redshift error at that redshift. Then, the number of independent triplets behind a cluster would be $N_{\text{bin}} - 2$, with N_{bin} the total number of bins. However, photometric redshifts, especially if some infrared filters are available, should be accurately determined.

In this paper, we have considered any halo mass profile but the method could be optimised by exploiting the symmetrical properties in the mass distribution of galaxy groups and clusters. A nearly elliptical matter distribution would allow to consider tangential shear averaged in concentric annuli, i.e. to collect triplets selecting galaxies in the concentric ring instead of a small local patch. This would make nearly sure that for any galaxy we can find a pair of galaxy redshifts that maximize the signal.

Some conclusions on the viability of the shear triplet method can be drawn by comparison with the shear ratio geometric test. The confidence regions we plotted seem larger than those obtained with the shear ratio test in a survey with similar properties (Taylor et al. 2007). As we have seen, the shear test is biased for large mass haloes $M \gtrsim 10^{15} M_\odot$, where the reduced shear should be properly considered instead of the shear. Being these haloes pretty rare if compared with halo masses of the order of $\lesssim 10^{14} M_\odot$, which provide the bulk of the signal, this systematic effect should not jeopardize the shear ratio method. In any case, since the two techniques require the same kind of measurements they should be properly integrated. This is desirable especially because the analysis of optically selected large mass halo in wide-field survey should begin with very massive haloes which, on turn, pay the highest dividend.

ACKNOWLEDGMENTS

MS is supported by the Swiss National Science Foundation and by the Tomalla Foundation.

REFERENCES

- Bartelmann M., Schneider P., 2001, *Phys. Rep.*, 340, 291
- Bernstein G., Jain B., 2004, *ApJ*, 600, 17
- Gautret L., Fort B., Mellier Y., 2000, *A&A*, 353, 10
- Golse G., Kneib J.-P., Soucail G., 2002, *A&A*, 387, 788
- Heavens A. F., Kitching T. D., Taylor A. N., 2006, *MNRAS*, 373, 105
- Hu W., Jain B., 2004, *Phys. Rev. D*, 70, 043009
- Huterer D., 2002, *Phys. Rev. D*, 65, 063001
- Jain B., Taylor A., 2003, *Phys. Rev. Lett.*, 91, 141302
- Linder E. V., 2004, *Phys. Rev. D*, 70, 043534
- Munshi D., Valageas P., Van Waerbeke L., Heavens A., 2006, preprint (astro-ph/0612667)

- Peacock J. A., Schneider P., Efstathiou G., Ellis J. R., Leibundgut B., Lilly S. J., Mellier Y., 2006, preprint (astro-ph/0610906)
- Seitz C., Schneider P., 1997, *A&A*, 318, 687
- Sereno M., 2002, *A&A*, 393, 757
- Sereno M., Longo G., 2004, *MNRAS*, 354, 1255
- Sereno M., Covone G., Piedipalumbo E., de Ritis R., 2001, *MNRAS*, 327, 517
- Sereno M., Piedipalumbo E., Sazhin M. V., 2002, *MNRAS*, 335, 1061
- Song Y.-S., Knox L., 2004, *Phys. Rev. D*, 70, 063510
- Takada M., Jain B., 2004, *MNRAS*, 348, 897
- Taylor A. N., Kitching T. D., Bacon D. J., Heavens A. F., 2007, *MNRAS*, 374, 1377
- Zhang J., Hui L., Stebbins A., 2005, *ApJ*, 635, 806

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.